

Beyond the spectral Standard Model: Pati-Salam unification

Walter van Suijlekom


(joint with Ali Chamseddine and Alain Connes)

22 September 2017


Radboud University Nijmegen



- Motivation: NCG and HEP
- Noncommutative Riemannian spin manifolds (aka spectral triples)
- Gauge theory from spectral triples: gauge group, gauge fields
- The spectral Standard Model and Beyond



A fermion in a spacetime background

- 
- Spacetime is a (pseudo) Riemannian manifold M :
local coordinates x_μ generate algebra $C^\infty(M)$.
 - Propagator is described by Dirac operator D_M , essentially a 'square root' of the Laplacian.

- The **Laplacian** on the circle \mathbb{S}^1 is given by

$$\Delta_{\mathbb{S}^1} = -\frac{d^2}{dt^2}; \quad (t \in [0, 2\pi))$$

- The **Dirac operator** on the circle is

$$D_{\mathbb{S}^1} = -i\frac{d}{dt}$$

with square $\Delta_{\mathbb{S}^1}$.



The 2-dimensional torus

- Consider the two-dimensional torus \mathbb{T}^2 parametrized by two angles $t_1, t_2 \in [0, 2\pi)$.
- The **Laplacian** reads

$$\Delta_{\mathbb{T}^2} = -\frac{\partial^2}{\partial t_1^2} - \frac{\partial^2}{\partial t_2^2}.$$

- At first sight it seems difficult to construct a differential operator that squares to $\Delta_{\mathbb{T}^2}$:

$$\left(a \frac{\partial}{\partial t_1} + b \frac{\partial}{\partial t_2} \right)^2 = a^2 \frac{\partial^2}{\partial t_1^2} + 2ab \frac{\partial^2}{\partial t_1 \partial t_2} + b^2 \frac{\partial^2}{\partial t_2^2}$$

- This puzzle was solved by Dirac who considered the possibility that a and b be complex *matrices*:

$$a = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}; \quad b = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$$

then $a^2 = b^2 = -1$ and $ab + ba = 0$

- The **Dirac operator on the torus** is

$$D_{\mathbb{T}^2} = \begin{pmatrix} 0 & \frac{\partial}{\partial t_1} + i \frac{\partial}{\partial t_2} \\ -\frac{\partial}{\partial t_1} + i \frac{\partial}{\partial t_2} & 0 \end{pmatrix},$$

which satisfies $(D_{\mathbb{T}^2})^2 = -\frac{\partial^2}{\partial t_1^2} - \frac{\partial^2}{\partial t_2^2}$.



The 4-dimensional torus

- Consider the 4-torus \mathbb{T}^4 parametrized by t_1, t_2, t_3, t_4 and the **Laplacian** is

$$\Delta_{\mathbb{T}^4} = -\frac{\partial^2}{\partial t_1^2} - \frac{\partial^2}{\partial t_2^2} - \frac{\partial^2}{\partial t_3^2} - \frac{\partial^2}{\partial t_4^2}.$$

- The search for a differential operator that squares to $\Delta_{\mathbb{T}^4}$ again involves matrices, but we also need **quaternions**:

$$i^2 = j^2 = k^2 = ijk = -1.$$

- The **Dirac operator** on \mathbb{T}^4 is

$$D_{\mathbb{T}^4} = \begin{pmatrix} 0 & \frac{\partial}{\partial t_1} + i \frac{\partial}{\partial t_2} + j \frac{\partial}{\partial t_3} + k \frac{\partial}{\partial t_4} \\ -\frac{\partial}{\partial t_1} + i \frac{\partial}{\partial t_2} + j \frac{\partial}{\partial t_3} + k \frac{\partial}{\partial t_4} & 0 \end{pmatrix}$$


- The relations $ij = -ji$, $ik = -ki$, *et cetera* imply that its square coincides with $\Delta_{\mathbb{T}^4}$.

Hearing the shape of a drum: motivation from math

Kac (1966), Connes (1989)

- The geometry of M is not fully determined by spectrum of D_M .
- This is considerably improved by considering besides D_M also the algebra $C^\infty(M)$ of smooth (coordinate) functions on M
- In fact, the Riemannian distance function on M is equal to

$$d(x, y) = \sup_{f \in C^\infty(M)} \{|f(x) - f(y)| : \text{gradient } f \leq 1\}$$



- The gradient of f is given by the commutator

$$[D_M, f] = D_M f - f D_M \quad (\text{e.g. } [D_{S^1}, f] = -i \frac{df}{dt})$$

Replace spacetime by **spacetime** \times **finite (nc) space**:
 $M \times F$

- F is considered as **internal space** (Kaluza–Klein like)
- F is described by a **noncommutative algebra**, such as $M_3(\mathbb{C})$, just as spacetime is described by **coordinate functions** $x_\mu(p)$.
- ‘Propagation’ of particles in F is described by a **Dirac-type operator** D_F which is actually simply a hermitian matrix.

- Finite space F , discrete topology

$$F = \quad 1 \bullet \quad 2 \bullet \quad \dots \quad N \bullet$$

- Smooth functions on F are given by N -tuples in \mathbb{C}^N , and the corresponding algebra $C^\infty(F)$ corresponds to **diagonal matrices**

$$\begin{pmatrix} f(1) & 0 & \dots & 0 \\ 0 & f(2) & \dots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \dots & f(N) \end{pmatrix}$$

- The **finite Dirac operator** is an arbitrary hermitian matrix D_F , giving rise to a distance function on F as

$$d(p, q) = \sup_{f \in C^\infty(F)} \{|f(p) - f(q)| : \|[D_F, f]\| \leq 1\}$$

Example: two-point space

$$F = 1 \bullet \quad 2 \bullet$$

- Then the **algebra of smooth functions**

$$C^\infty(F) := \left\{ \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \mid \lambda_1, \lambda_2 \in \mathbb{C} \right\}$$

- A **finite Dirac operator** is given by

$$D_F = \begin{pmatrix} 0 & \bar{c} \\ c & 0 \end{pmatrix}; \quad (c \in \mathbb{C})$$

- The **distance formula** then becomes

$$d(1, 2) = \frac{1}{|c|}$$



The geometry of F gets much more interesting if we allow for a *noncommutative* structure at each point of F .

- Instead of diagonal matrices, we consider **block diagonal** matrices

$$A = \begin{pmatrix} a_1 & 0 & \cdots & 0 \\ 0 & a_2 & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & a_N \end{pmatrix},$$

where the a_1, a_2, \dots, a_N are square matrices of size n_1, n_2, \dots, n_N .

- Hence we will consider the **matrix algebra**

$$\mathcal{A}_F := M_{n_1}(\mathbb{C}) \oplus M_{n_2}(\mathbb{C}) \oplus \cdots \oplus M_{n_N}(\mathbb{C})$$

- A **finite Dirac operator** is still given by a hermitian matrix.

Example: noncommutative two-point space

The two-point space can be given a noncommutative structure by considering the algebra \mathcal{A}_F of 3×3 block diagonal matrices of the following form

$$\begin{pmatrix} \lambda & 0 & 0 \\ 0 & a_{11} & a_{12} \\ 0 & a_{21} & a_{22} \end{pmatrix}$$

A finite Dirac operator for this example is given by a hermitian 3×3 matrix, for example

$$D_F = \begin{pmatrix} 0 & \bar{c} & 0 \\ c & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$



$$(\mathcal{A}, \mathcal{H}, D)$$

- Extended to **real** spectral triple:
 - $J : \mathcal{H} \rightarrow \mathcal{H}$ real structure (**charge conjugation**)
- such that

$$J^2 = \pm 1; \quad JD = \pm DJ$$

- **Right action of \mathcal{A}** on \mathcal{H} : $a^{\text{op}} = Ja^*J^{-1}$ so that $(ab)^{\text{op}} = b^{\text{op}}a^{\text{op}}$ and

$$[a^{\text{op}}, b] = 0; \quad a, b \in \mathcal{A}$$

- D is said to satisfy **first-order condition** if

$$[[D, a], b^{\text{op}}] = 0$$



Spectral invariants

Chamseddine–Connes (1996, 1997)

$$\text{Trace } f(D/\Lambda) + \frac{1}{2} \langle J\tilde{\psi}, D\tilde{\psi} \rangle$$

- **Invariant** under unitaries $u \in \mathcal{U}(\mathcal{A})$ acting as

$$D \mapsto UDU^*; \quad U = u(u^*)^{\text{op}}$$

- **Gauge group**: $\mathcal{G}(\mathcal{A}) := \{u(u^*)^{\text{op}} : u \in \mathcal{U}(\mathcal{A})\}$.
- Compute *rhs*:

$$D \mapsto D + u[D, u^*] \pm Ju[D, u^*]J^{-1}$$



Semigroup of inner perturbations

Chamseddine–Connes-vS (2013)

Extend this to more general perturbations:

$$\text{Pert}(\mathcal{A}) := \left\{ \sum_j a_j \otimes b_j^{\text{op}} \in \mathcal{A} \otimes \mathcal{A}^{\text{op}} \left| \begin{array}{l} \sum_j a_j b_j = 1 \\ \sum_j a_j \otimes b_j^{\text{op}} = \sum_j b_j^* \otimes a_j^{\text{op}} \end{array} \right. \right\}$$

with semi-group law inherited from product in $\mathcal{A} \otimes \mathcal{A}^{\text{op}}$.

- $\mathcal{U}(\mathcal{A})$ maps to $\text{Pert}(\mathcal{A})$ by sending $u \mapsto u \otimes u^{\text{op}}$.
- $\text{Pert}(\mathcal{A})$ acts on D :

$$D \mapsto \sum_j a_j D b_j = D + \sum_j a_j [D, b_j]$$

and this also extends to real spectral triples via the map

$$\text{Pert}(\mathcal{A}) \rightarrow \text{Pert}(\mathcal{A} \otimes JAJ^{-1})$$

Proposition

Let \mathcal{A}_F be the algebra of block diagonal matrices (fixed size).
Then the *perturbation semigroup* of \mathcal{A}_F is

$$\text{Pert}(\mathcal{A}_F) \simeq \left\{ \sum_j A_j \otimes B_j \in \mathcal{A}_F \otimes \mathcal{A}_F \left| \begin{array}{l} \sum_j A_j (B_j)^t = \mathbb{I} \\ \sum_j A_j \otimes B_j = \sum_j \bar{B}_j \otimes \bar{A}_j \end{array} \right. \right\}$$

The semigroup law in $\text{Pert}(\mathcal{A}_F)$ is given by the matrix product in $\mathcal{A}_F \otimes \mathcal{A}_F$:

$$(A \otimes B)(A' \otimes B') = (AA') \otimes (BB').$$

Example: perturbation semigroup of two-point space

- Now $\mathcal{A}_F = \mathbb{C}^2$, the algebra of diagonal 2×2 matrices.
- In terms of the standard basis of such matrices

$$e_{11} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad e_{22} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

we can write an arbitrary element of $\text{Pert}(\mathbb{C}^2)$ as

$$z_1 e_{11} \otimes e_{11} + z_2 e_{11} \otimes e_{22} + z_3 e_{22} \otimes e_{11} + z_4 e_{22} \otimes e_{22}$$

- Matrix multiplying e_{11} and e_{22} yields for the normalization condition:

$$z_1 = 1 = z_4.$$

- The self-adjointness condition reads

$$z_2 = \bar{z}_3$$

leaving only one free complex parameter so that $\text{Pert}(\mathbb{C}^2) \simeq \mathbb{C}$.

- More generally, $\text{Pert}(\mathbb{C}^N) \simeq \mathbb{C}^{N(N-1)/2}$ with componentwise product.

Example: perturbation semigroup of $M_2(\mathbb{C})$

- Let us consider a **noncommutative example**, $\mathcal{A}_F = M_2(\mathbb{C})$.
- We can identify $M_2(\mathbb{C}) \otimes M_2(\mathbb{C})$ with $M_4(\mathbb{C})$ so that elements in $\text{Pert}(M_2(\mathbb{C}))$ are **4×4 -matrices** satisfying the **normalization** and **self-adjointness** condition. In a suitable basis:

$$\text{Pert}(M_2(\mathbb{C})) = \left\{ \begin{array}{c} \begin{pmatrix} 1 & v_1 & v_2 & iv_3 \\ 0 & x_1 & x_2 & ix_3 \\ 0 & x_4 & x_5 & ix_6 \\ 0 & ix_7 & ix_8 & x_9 \end{pmatrix} \mid \begin{array}{l} v_1, v_2, v_3 \in \mathbb{R} \\ x_1, \dots, x_9 \in \mathbb{R} \end{array} \end{array} \right\}$$

and one can show that

$$\text{Pert}(M_2(\mathbb{C})) \simeq \mathbb{R}^3 \rtimes S.$$

- More generally (B.Sc. thesis Niels Neumann),

$$\text{Pert}(M_N(\mathbb{C})) \simeq W \rtimes S'.$$

Example: noncommutative two-point space

- Consider **noncommutative two-point space** described by $\mathbb{C} \oplus M_2(\mathbb{C})$
- It turns out that

$$\text{Pert}(\mathbb{C} \oplus M_2(\mathbb{C})) \simeq M_2(\mathbb{C}) \times \text{Pert}(M_2(\mathbb{C}))$$

- Only $M_2(\mathbb{C}) \subset \text{Pert}(\mathbb{C} \oplus M_2(\mathbb{C}))$ acts non-trivially on D_F :

$$D_F = \begin{pmatrix} 0 & \bar{c} & 0 \\ c & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \mapsto \begin{pmatrix} 0 & \bar{c}\bar{\phi}_1 & \bar{c}\bar{\phi}_2 \\ c\phi_1 & 0 & 0 \\ c\phi_2 & 0 & 0 \end{pmatrix}$$

- Physicists call ϕ_1 and ϕ_2 the **Higgs field**.
- The **group of unitary block diagonal matrices** is now $U(1) \times U(2)$ and an element (λ, u) therein acts as

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix} \mapsto \bar{\lambda} u \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}.$$

Example: perturbation semigroup of a manifold

Recall, for any involutive algebra \mathcal{A}

$$\text{Pert}(\mathcal{A}) := \left\{ \sum_j a_j \otimes b_j^{\text{op}} \in \mathcal{A} \otimes \mathcal{A}^{\text{op}} \left| \begin{array}{l} \sum_j a_j b_j = 1 \\ \sum_j a_j \otimes b_j^{\text{op}} = \sum_j b_j^* \otimes a_j^{*\text{op}} \end{array} \right. \right\}$$

- We can consider functions in $C^\infty(M) \otimes C^\infty(M)$ as functions of two variables in $C^\infty(M \times M)$.
- The normalization and self-adjointness condition in $\text{Pert}(C^\infty(M))$ translate accordingly and yield

$$\text{Pert}(C^\infty(M)) = \left\{ f \in C^\infty(M \times M) \left| \begin{array}{l} f(x, x) = 1 \\ f(x, y) = \overline{f(y, x)} \end{array} \right. \right\}$$


- The action of $\text{Pert}(C^\infty(M))$ on the partial derivatives appearing in a **Dirac operator** D_M is given by

$$\frac{\partial}{\partial x_\mu} \mapsto \frac{\partial}{\partial x_\mu} + \frac{\partial}{\partial y_\mu} f(x, y) \Big|_{y=x} =: \partial_\mu + A_\mu$$

- Combine (4d) Riemannian spin manifold M with finite noncommutative space F :

$$M \times F$$

- F is internal space at each point of M



- Described by matrix-valued functions on M : algebra $C^\infty(M, \mathcal{A}_F)$



Dirac operator on $M \times F$

- Recall the form of D_M :

$$D_M = \begin{pmatrix} 0 & D_M^+ \\ D_M^- & 0 \end{pmatrix}.$$

- Dirac operator on $M \times F$ is the combination

$$D_{M \times F} = D_M + \gamma_5 D_F = \begin{pmatrix} D_F & D_M^+ \\ D_M^- & -D_F \end{pmatrix}.$$

- The crucial property of this specific form is that it squares to the sum of the two Laplacians on M and F :

$$D_{M \times F}^2 = D_M^2 + D_F^2$$

- Using this, we can expand the heat trace:


$$\text{Trace } e^{-D_{M \times F}^2 / \Lambda^2} = \frac{\text{Vol}(M) \Lambda^4}{(4\pi)^2} \text{Trace} \left(1 - \frac{D_F^2}{\Lambda^2} + \frac{D_F^4}{2\Lambda^4} \right) + \mathcal{O}(\Lambda^{-1}).$$

The Higgs mechanism

We apply this to the noncommutative two-point space described before

- Algebra $\mathcal{A}_F = \mathbb{C} \oplus M_2(\mathbb{C})$
- **Perturbation** of Dirac operator D_M parametrized by gauge bosons for $U(1) \times U(2)$.
- **Perturbation** of finite Dirac operator D_F parametrized by ϕ_1, ϕ_2 .
- Spectral action for the perturbed Dirac operator induces a potential:

$$V(\phi) = -2\Lambda^2(|\phi_1|^2 + |\phi_2|^2) + (|\phi_1|^2 + |\phi_2|^2)^2$$



The spectral Standard Model

Describe $M \times F_{SM}$ by [CCM 2007]

- **Coordinates:** $\hat{x}^\mu(p) \in \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ (with unimodular unitaries $U(1)_Y \times SU(2)_L \times SU(3)$).
- **Dirac operator** $D_{M \times F} = D_M + \gamma_5 D_F$ where

$$D_F = \begin{pmatrix} S & T^* \\ T & \bar{S} \end{pmatrix}$$

is a 96×96 -dimensional hermitian matrix where 96 is:

$$3 \times 2 \times (\underbrace{2}_{\substack{\uparrow \\ \text{families}}} \otimes \underbrace{1}_{\substack{\uparrow \\ \text{anti-particles}}} + \underbrace{1}_{\nu_R} \otimes \underbrace{1}_{e_R} + \underbrace{1}_{e_R} \otimes \underbrace{1}_{(u_L, d_L)} + \underbrace{2}_{(u_L, d_L)} \otimes \underbrace{3}_{u_R} + \underbrace{1}_{u_R} \otimes \underbrace{3}_{d_R} + \underbrace{1}_{d_R} \otimes \underbrace{3}_{d_R})$$

$$D_F = \begin{pmatrix} S & T^* \\ T & \bar{S} \end{pmatrix}$$

- The operator S is given by

$$S_l := \begin{pmatrix} 0 & 0 & Y_\nu & 0 \\ 0 & 0 & 0 & Y_e \\ Y_\nu^* & 0 & 0 & 0 \\ 0 & Y_e^* & 0 & 0 \end{pmatrix}, \quad S_q \otimes 1_3 = \begin{pmatrix} 0 & 0 & Y_u & 0 \\ 0 & 0 & 0 & Y_d \\ Y_u^* & 0 & 0 & 0 \\ 0 & Y_d^* & 0 & 0 \end{pmatrix} \otimes 1_3,$$

where Y_ν , Y_e , Y_u and Y_d are 3×3 mass matrices acting on the three generations.

- The symmetric operator T only acts on the right-handed (anti)neutrinos, $T\nu_R = Y_R\bar{\nu}_R$ for a 3×3 symmetric Majorana mass matrix Y_R , and $Tf = 0$ for all other fermions $f \neq \nu_R$.

- Inner perturbations of D_M give a matrix

$$A_\mu = \begin{pmatrix} B_\mu & 0 & 0 & 0 \\ 0 & W_\mu^3 & W_\mu^+ & 0 \\ 0 & W_\mu^- & -W_\mu^3 & 0 \\ 0 & 0 & 0 & (G_\mu^a) \end{pmatrix}$$

corresponding to **hypercharge, weak and strong interaction**.

- Inner perturbations of D_F give

$$\begin{pmatrix} Y_\nu & 0 \\ 0 & Y_e \end{pmatrix} \rightsquigarrow \begin{pmatrix} Y_\nu \phi_1 & -Y_e \bar{\phi}_2 \\ Y_\nu \phi_2 & Y_e \bar{\phi}_1 \end{pmatrix}$$

corresponding to **SM-Higgs field**. Similarly for Y_u, Y_d .

If we consider the spectral action:

$$\text{Trace } f(D_M/\Lambda) \sim c_0 \int F_{\mu\nu} F^{\mu\nu} - c'_2 |\phi|^2 + c'_0 |\phi|^4 + \dots$$

we observe [CCM 2007]:

- The coupling constants of hypercharge, weak and strong interaction are expressed in terms of the **single constant** c_0 which implies

$$g_3^2 = g_2^2 = \frac{5}{3} g_1^2$$

In other words, there should be **grand unification**.

- Moreover, the quartic Higgs coupling λ is related via


$$\lambda \approx 24 \frac{3 + \rho^4}{(3 + \rho^2)^2} g_2^2; \quad \rho = \frac{m_\nu}{m_{\text{top}}}$$

Phenomenology of the spectral Standard Model

This can be used to derive predictions as follows:

- Interpret the spectral action as an **effective field theory** at $\Lambda_{\text{GUT}} \approx 10^{13} - 10^{16}$ GeV.
- Run the quartic coupling constant λ to SM-energies to predict

$$m_h^2 = \frac{4\lambda M_W^2}{3g_2^2}$$





This gives [CCM 2007]

$$167 \text{ GeV} \leq m_h \leq 176 \text{ GeV}$$

Three problems

- 1 This prediction is **falsified** by the now measured value.
- 2 In the Standard Model there is not the **presumed grand unification**.
- 3 There is a problem with the low value of m_h , making the Higgs vacuum un/metastable [Elias-Miro et al. 2011].



Beyond the SM with noncommutative geometry

A solution to the above three problems?

- The matrix coordinates of the Standard Model arise naturally as a restriction of the following **coordinates**

$$\hat{x}^\mu(p) = (q_R^\mu(p), q_L^\mu(p), m^\mu(p)) \in \mathbb{H}_R \oplus \mathbb{H}_L \oplus M_4(\mathbb{C})$$

corresponding to a **Pati–Salam unification**:

$$U(1)_Y \times SU(2)_L \times SU(3) \rightarrow SU(2)_R \times SU(2)_L \times SU(4)$$

- The 96 **fermionic degrees of freedom** are structured as

$$\left(\begin{array}{cc|cc} \nu_R & u_{iR} & \nu_L & u_{iL} \\ e_R & d_{iR} & e_L & d_{iL} \end{array} \right) \quad (i = 1, 2, 3)$$


- Again the **finite Dirac operator** is a 96×96 -dimensional matrix (details in [CCS 2013]).

- Inner perturbations of D_M now give **three gauge bosons**:

$$W_R^\mu, \quad W_L^\mu, \quad V^\mu$$

corresponding to $SU(2)_R \times SU(2)_L \times SU(4)$.

- For the inner perturbations of D_F we distinguish two cases, depending on the initial form of D_F :
 - I The Standard Model $D_F = \begin{pmatrix} S & T^* \\ T & \bar{S} \end{pmatrix}$
 - II A more general D_F with zero $\bar{f}_L - f_L$ -interactions.



Scalar sector of the spectral Pati–Salam model

Case I For a SM D_F , the resulting scalar fields are **composite fields**, expressed in scalar fields whose representations are:

	$SU(2)_R$	$SU(2)_L$	$SU(4)$
$\phi_{\dot{a}}^b$	2	2	1
$\Delta_{\dot{a}I}$	2	1	4
Σ_J^I	1	1	15

Case II For a more general finite Dirac operator, we have **fundamental scalar fields**:


particle	$SU(2)_R$	$SU(2)_L$	$SU(4)$
$\Sigma_{\dot{a}J}^{bJ}$	2	2	1 + 15
$H_{\dot{a}I}^{bJ} \left\{ \right.$	3	1	10
	1	1	6

As for the Standard Model, we can compute the spectral action which describes the usual **Pati–Salam model** with

- **unification** of the gauge couplings

$$g_R = g_L = g.$$

- A rather involved, fixed **scalar potential**, still subject to further study



However, independently from the spectral action, we can analyze the running at one loop of the gauge couplings [CCS 2015]:

- 1 We run the **Standard Model gauge couplings** up to a presumed PS \rightarrow SM symmetry breaking scale m_R
- 2 We take their values as **boundary conditions** to the **Pati–Salam gauge couplings** g_R, g_L, g at this scale via

$$\frac{1}{g_1^2} = \frac{2}{3} \frac{1}{g^2} + \frac{1}{g_R^2}, \quad \frac{1}{g_2^2} = \frac{1}{g_L^2}, \quad \frac{1}{g_3^2} = \frac{1}{g^2},$$

- 3 Vary m_R in a search for a **unification scale** Λ where


$$g_R = g_L = g$$

which is where the **spectral action** is valid as an **effective theory**.

Phenomenology of the spectral Pati–Salam model

Case I: Standard Model D_F

For the **Standard Model Dirac operator**, we have found that with $m_R \approx 4.25 \times 10^{13}$ GeV there is unification at $\Lambda \approx 2.5 \times 10^{15}$ GeV:



Phenomenology of the spectral Pati–Salam model

Case I: Standard Model D_F

In this case, we can also say something about the **scalar particles** that remain after SSB:

	$U(1)_Y$	$SU(2)_L$	$SU(3)$
$\begin{pmatrix} \phi_1^0 \\ \phi_1^+ \end{pmatrix} = \begin{pmatrix} \phi_1^1 \\ \phi_1^2 \end{pmatrix}$	1	2	1
$\begin{pmatrix} \phi_2^- \\ \phi_2^0 \end{pmatrix} = \begin{pmatrix} \phi_2^1 \\ \phi_2^2 \end{pmatrix}$	-1	2	1
σ	0	1	1
η	$-\frac{2}{3}$	1	3


- It turns out that these scalar fields have a **little influence** on the running of the SM-gauge couplings (at one loop).
- However, this sector contains the **real scalar singlet** σ that allowed for a **realistic Higgs mass** and that **stabilizes** the Higgs vacuum [CC 2012].

Phenomenology of the spectral Pati–Salam model

Case II: General Dirac

For the more general case, we have found that with

$m_R \approx 1.5 \times 10^{11}$ GeV there is **unification** at $\Lambda \approx 6.3 \times 10^{16}$ GeV:



Conclusion

We have arrived at a **spectral Pati–Salam model** that

- goes beyond the Standard Model
- has a **fixed scalar sector** once the finite Dirac operator has been fixed (only a **few scenarios**)
- exhibits **grand unification** for all of these scenarios (confirmed by [Aydemir–Minic–Sun–Takeuchi 2015])
- the scalar sector has the potential to **stabilize the Higgs vacuum** and allow for a **realistic Higgs mass**.

A. Chamseddine, A. Connes, WvS.

Beyond the Spectral Standard Model: Emergence of
Pati-Salam Unification. *JHEP* 11 (2013) 132.
[arXiv:1304.8050]

Grand Unification in the Spectral Pati-Salam Model. *JHEP* 11
(2015) 011. [arXiv:1507.08161]

WvS.

Noncommutative Geometry and Particle Physics.
Mathematical Physics Studies, Springer, 2015.

and also: <http://www.noncommutativegeometry.nl>