

# Operator systems and noncommutative geometry

Walter van Suijlekom

Radboud Universiteit



# The spectral approach to geometry


Given cpt Riemannian spin manifold  $(M, g)$  with spinor bundle  $S_M$  on  $M$ .

- ▶ the  $C^*$ -algebra  $C(M)$
- ▶ the self-adjoint Dirac operator  $D_M$
- ▶ both acting on Hilbert space  $L^2(S_M)$

$\rightsquigarrow$  spectral triple:  $(C(M), L^2(S_M), D_M)$

Reconstruction of distance function [Connes 1994]:

$$d(x, y) = \sup_{f \in C(M)} \{ |\text{ev}_x(f) - \text{ev}_y(f)| : \|[D_M, f]\| \leq 1 \}$$



# Spectral triples

More generally, we consider a triple  $(A, \mathcal{H}, D)$

- ▶ a  $C^*$ -algebra  $A$
- ▶ a self-adjoint operator  $D$  with compact resolvent and bounded commutators  $[D, a]$  for  $a \in \mathcal{A} \subset A$
- ▶ both acting (boundedly, resp. unboundedly) on Hilbert space  $\mathcal{H}$

Generalized distance function:

- ▶ States are positive linear functionals  $\phi : A \rightarrow \mathbb{C}$  of norm 1
- ▶ Pure states are extreme points of state space
- ▶ Distance function on state space of  $A$ :

$$d(\phi, \psi) = \sup_{a \in A} \{ |\phi(a) - \psi(a)| : \|[D, a]\| \leq 1 \}$$

## Spectral data: $(\mathcal{A}, \mathcal{H}, D)$

- ▶ The mathematical reformulation of geometry in terms of spectral data (global analysis) requires the knowledge of the full Dirac operator.
- ▶ From a physical standpoint this is not very realistic: detectors have limited energy ranges and resolution.
- ▶ We aim for the underlying mathematical formalism for doing (noncommutative) geometry with only part of the spectrum and/or with finite resolution.

This is in line with [D'Andrea–Lizzi–Martinetti 2014], [Glaser–Stern 2019] and based on [Connes–vS] (CMP, Szeged)

# Operator systems

(I) Given  $(A, \mathcal{H}, D)$  we project onto part of the spectrum of  $D$ :

- ▶  $\mathcal{H} \mapsto P\mathcal{H}$ , projection onto closed Hilbert subspace
- ▶  $D \mapsto PDP$ , still a self-adjoint operator
- ▶  $A \mapsto PAP$ , this is not an algebra any more (unless  $P \in A$ )

Instead,  $PAP$  is an operator system:  $(PaP)^* = Pa^*P$ .

(II) Another approach would be to consider metric spaces up to a finite resolution :

- ▶ Consider integral operators associated to the tolerance relation  $R_\epsilon$  given by  $d(x, y) < \epsilon$

So first, some background on operator systems.

# Operator systems

Definition (Arveson 1969, Choi-Effros 1977)

*An operator system is a  $*$ -closed vector space  $E$  of bounded operators. Unital: it contains the identity operator.*

- ▶  $E$  is ordered: cone  $E_+ \subseteq E$  of positive operators, in the sense that  $T \in E_+$  iff

$$\langle \psi, T\psi \rangle \geq 0; \quad (\psi \in \mathcal{H}).$$

- ▶ in fact,  $E$  is matrix ordered: cones  $M_n(E)_+ \subseteq M_n(E)$  of positive operators on  $\mathcal{H}^n$  for any  $n$ .

Maps between operator systems  $E, F$  are completely positive maps in the sense that their extensions  $M_n(E) \rightarrow M_n(F)$  are positive for all  $n$ .

Isomorphisms are complete order isomorphisms


# $C^*$ -envelope of a unital operator system

Arveson introduced the notion of  $C^*$ -envelope for unital operator systems in 1969, Hamana established existence and uniqueness in 1979. Non-unital case: [Connes-vS 2020], [Kennedy–Kim–Manor 2021]

A  $C^*$ -extension  $\kappa : E \rightarrow A$  of a unital operator system  $E$  is given by a complete order isomorphism onto  $\kappa(E) \subseteq A$  such that  $C^*(\kappa(E)) = A$ .

A  $C^*$ -envelope of a unital operator system is a  $C^*$ -extension  $\kappa : E \rightarrow A$  with the following universal property:

$$\begin{array}{ccc} E & \xrightarrow{\kappa} & A \\ & \searrow \lambda & \uparrow \exists! \rho \\ & & B \end{array}$$



# Shilov boundaries

There is a useful description of  $C^*$ -envelopes using Shilov ideals.

## Definition

Let  $\kappa : E \rightarrow A$  be a  $C^*$ -extension of an operator system. A boundary ideal is given by a closed 2-sided ideal  $I \subseteq A$  such that the quotient map  $q : A \rightarrow A/I$  is completely isometric on  $\kappa(E) \subseteq A$ .

*The Shilov boundary ideal is the largest of such boundary ideals.*

## Proposition (Hamana 1979)

Let  $\kappa : E \rightarrow A$  be a  $C^*$ -extension. Then there exists a Shilov boundary ideal  $J$  and  $C_{env}^*(E) \cong A/J$ .

As an example consider the operator system of continuous harmonic functions  $C_{\text{harm}}(\overline{\mathbb{D}})$  on the closed disc. Then by the maximum modulus principle the Shilov boundary is  $S^1$ . Accordingly, its  $C^*$ -envelope is  $C(S^1)$ .



# $C^*$ -envelope via boundary representations (à la Arveson)

Dritschel–McCullough 2005, Arveson 2008, Davidson–Kennedy 2015

View  $E$  as a concrete operator system in the  $C^*$ -algebra  $C^*(E)$  it generates.

- ▶ A ucp map  $\phi : E \rightarrow B(\mathcal{H})$  has the *unique extension property* if it has a unique ucp extension to  $C^*(E)$  which is a  $*$ -representation
- ▶ If, in addition, the  $*$ -representation is irreducible, it is called a *boundary representation*

## Theorem

*A unital operator system  $E$  is completely normed by its boundary representations, that is, for every  $x \in M_n(E)$  there is a boundary representation  $\pi$  of  $E$  such that  $\|x\| = \|\pi^{(n)}(x)\|$ .*

## Corollary

*The direct sum of all boundary representations of  $E$  yield a completely isometric map  $\iota : E \rightarrow B(\mathcal{H})$  such that  $(C^*(\iota(E)), \iota)$  is the  $C^*$ -envelope of  $E$ .*

# Propagation number of an operator system

One lets  $E^{\circ n}$  be the norm closure of the linear span of products of  $\leq n$  elements of  $E$ .

## Definition

*The propagation number  $\text{prop}(E)$  of  $E$  is defined as the smallest integer  $n$  such that  $\kappa(E)^{\circ n} \subseteq C_{\text{env}}^*(E)$  is a  $C^*$ -algebra.*

Returning to harmonic functions in the disk we have  $\text{prop}(C_{\text{harm}}(\overline{\mathbb{D}})) = 1$ .

Proposition (Connes-vS, 2020; Pawłowska, 2024)

*The propagation number is invariant under complete order isomorphisms, as well as under stable=Morita equivalence [EKT, 2019]:*

$$\text{prop}(E) = \text{prop}(E \otimes_{\min} \mathcal{K})$$

*More generally [Koot, 2021], we have*

$$\text{prop}(E \otimes_{\min} F) = \max\{\text{prop}(E), \text{prop}(F)\}$$

# Spectral truncation of the circle [Connes-vS, 2020]

- ▶ Eigenvectors of  $D_{S^1}$  are Fourier modes  $e_k(t) = e^{ikt}$  for  $k \in \mathbb{Z}$
- ▶ Orthogonal projection  $P = P_n$  onto  $\text{span}_{\mathbb{C}}\{e_1, e_2, \dots, e_n\}$
- ▶ The space  $C(S^1)^{(n)} := PC(S^1)P$  is an operator system
- ▶ Any  $T = PfP$  in  $C(S^1)^{(n)}$  can be written as a Toeplitz matrix

$$PfP \sim (t_{k-l})_{kl} = \begin{pmatrix} t_0 & t_{-1} & \cdots & t_{-n+2} & t_{-n+1} \\ t_1 & t_0 & t_{-1} & & t_{-n+2} \\ \vdots & t_1 & t_0 & \ddots & \vdots \\ & & \ddots & \ddots & t_{-1} \\ t_{n-2} & & & & t_{-1} \\ t_{n-1} & t_{n-2} & \cdots & t_1 & t_0 \end{pmatrix}$$

We have:  $C_{\text{env}}^*(C(S^1)^{(n)}) \cong M_n(\mathbb{C})$

# Dual operator system: Fejér–Riesz

We introduce the Fejér–Riesz operator system  $C(S^1)_{(n)}$ :

- ▶ functions on  $S^1$  with a finite number of non-zero Fourier coefficients:

$$a = (\dots, 0, a_{-n+1}, a_{-n+2}, \dots, a_{-1}, a_0, a_1, \dots, a_{n-2}, a_{n-1}, 0, \dots)$$

- ▶ an element  $a$  is positive iff  $\sum_k a_k e^{ikx}$  is a positive function on  $S^1$ .
- ▶ The  $C^*$ -envelope of  $C(S^1)_{(n)}$  is given by  $C^*(\mathbb{Z})$

## Proposition

1. *The extreme rays in  $(C(S^1)_{(n)})_+$  are given by the elements  $a = (a_k)$  for which the Laurent series  $\sum_k a_k z^k$  has all its zeroes on  $S^1$ .*
2. *The pure states of  $C(S^1)_{(n)}$  are given by  $a \mapsto \sum_k a_k \lambda^k$  ( $\lambda \in S^1$ ).*

# Pure states on the Toeplitz matrices

Duality of  $C(S^1)^{(n)}$  and  $C(S^1)_{(n)}$  [Connes–vS 2020] and [Farenick 2021]:

$$\begin{aligned} C(S^1)^{(n)} \times C(S^1)_{(n)} &\rightarrow \mathbb{C} \\ (T = (t_{k-l})_{k,l}, a = (a_k)) &\mapsto \sum_k a_k t_{-k} \end{aligned}$$

## Proposition

1. The extreme rays in  $C(S^1)_+^{(n)}$  are  $\gamma(\lambda) = |f_\lambda\rangle\langle f_\lambda|$  for any  $\lambda \in S^1$ .
2. The pure state space  $\mathcal{P}(C(S^1)^{(n+1)}) \cong \mathbb{T}^n/S_n$ .

# Curiosities on Toeplitz matrices

Theorem (Carathéodory)

Let  $T$  be an  $n \times n$  Toeplitz matrix. Then  $T \geq 0$  iff  $T = V\Delta V^*$  with

$$\Delta = \begin{pmatrix} d_1 & & & \\ & d_2 & & \\ & & \dots & \\ & & & d_n \end{pmatrix}; \quad V = \frac{1}{\sqrt{n}} \begin{pmatrix} 1 & 1 & \dots & 1 \\ \lambda_1 & \lambda_2 & \dots & \lambda_n \\ \vdots & & & \vdots \\ \lambda_1^{n-1} & \lambda_2^{n-1} & \dots & \lambda_n^{n-1} \end{pmatrix},$$

for some  $d_1, \dots, d_n \geq 0$  and  $\lambda_1, \dots, \lambda_n \in S^1$ .